Prescribing Information

Presentation
White egg-shaped
pessaries each
containing 150mg
econazole nitrate and
melting at 37°C.

lises

Vaginitis due to Candida albicans and other veasts.

Dosage and Administration

One pessary should be inserted high into the vagina each evening for three consecutive days.

Contra-indications, Warnings etc.

Contra-indications:
None known.
Side-effects:
None known.
Overdosage:
Not applicable.
Precautions:
Hypersensitivity has

nypersensitivity has never been recorded, but if it should occur, administration of the product should be discontinued.

Gyno-Pevaryl*150
vaginal pessaries are
effective in the
treatment of vaginal
candidiasis associated
with pregnancy but as
with any new product,
caution should be
exercised when
prescribing during the

first trimester.

Pharmaceutical

Precautions

Store in a cool place.

Legal Category

Prescription only (POM).
Package Quantities
Each pack contains
3 pessaries and a patient
instruction leaflet.
The pessaries are
individually sealed in
a white plastic strip
marked Gyno-Pevary!*


150 at regular intervals. Further Information Anogenital hygiene is important to help prevent re-infection.

PL0076/0058
Name and address

of Licensee Ortho Pharmaceutical Limited, P.O. Box 79, Saunderton, High Wycombe, Buckinghamshire HP14 4HJ, England.

Further information and data sheet available on request.





Rapid symptomatic relief Burning, pruritus and discharge promptly subside on therapy.

Effective in over 85% of cases—in only 3 days! Three days are all it takes to clear the condition in over 85% of cases.!

Better patient compliance Three rapidly dissolving soft-wax pessaries inserted one at night for three days, effectively minimise the possibility of recurrence through non-compliance by reducing the therapy to its simplest form ever. Gyno-Pevaryl*150 vaginal pessaries – easy, quick and effective in the treatment of vaginal candidiasis.

Gyño-Pevaryl*150

Vaginal Pessaries

Econazole Nitrate 150mg.

new-generation antifungal

the effective 3 day, 3 pessary treatment

*Trademark © OPL 1978

Reference: 1. Balmer, J.A. (1976). Am J. Obstet. Gynae. 126, 4, 436-441.